Deformed Richardson-Gaudin model

P. Kulish (1), A. Stolin (2), and H. Johannesson (3)

(1) Steklov Mathematical Institute, 27, Fontanka, 191023 St. Petersburg, Russia
(2) Department of Mathematics, Chalmers University of Technology and University of Gothenburg, SE 412 96 Gothenburg, Sweden
(3) Department of Physics, University of Gothenburg, SE 412 96 Gothenburg, Sweden

arXiv:1402.4838v1 [nlin.SI] 19 Feb 2014

Abstract. The Richardson-Gaudin model describes strong pairing correlations of fermions confined to a finite chain. The integrability of the Hamiltonian allows for its eigenstates to be constructed algebraically. In this work we show that quantum group theory provides a possibility to deform the Hamiltonian while preserving integrability. More precisely, we use the so-called Jordanian r-matrix to deform the Hamiltonian of the Richardson-Gaudin model. In order to preserve its integrability, we need to insert a special nilpotent term into the auxiliary L-operator which generates integrals of motion of the system. Moreover, the quantum inverse scattering method enables us to construct the exact eigenstates of the deformed Hamiltonian. These states have a highly complex entanglement structure which requires further investigation.

PACS numbers: 02.30.Ik, 03.65.Fd, 21.10.Re, 74.20.Rp

The Richardson-Gaudin model [1, 2] is an integrable spin-1/2 periodic chain with Hamiltonian

$$H = \sum_{j=1}^{N} \epsilon_j S_j^z + g \sum_{j,\,k=1}^{N} S_j^- S_k^+, \tag{1}$$

where g is a coupling constant and $S_l^{\pm} = (S_l^x \pm i S_l^y)$, with S_l^α N copies of the Lie algebra $su(2)$ generators

$$[S_l^\alpha, S_{l'}^\beta] = i\varepsilon^{\alpha\,\beta\,\gamma} S^\gamma \delta_{l\,l'}, \qquad \alpha, \beta, \gamma = x, y, z. \tag{2}$$

As shown by Cambiaggio *et al.* [3], by introducing fermion operators $c_{lm}^\dagger$ and c_{lm} related to the sl(2) generators by

$$S_l^z = \frac{1}{2}\sum_m c_{lm}^\dagger c_{lm} - \frac{1}{2}, \quad S_l^+ = \frac{1}{2}\sum_m c_{lm}^\dagger c_{l\bar{m}}^\dagger = (S_l^-)^\dagger, \tag{3}$$

the Richardson-Gaudin model in Eq. (1) gets mapped onto the pairing model Hamiltonian

$$H_P = \sum_l \epsilon_l \hat{n}_l + \frac{g}{2}\sum_{l,l'} A_l^\dagger A_{l'}. \tag{4}$$

Here $c_{lm}^\dagger$ (c_{lm}) creates (annihilates) a fermion in the state $|\,lm\rangle$ (with $|\,l\bar{m}\rangle$ the time reversed state of $|\,lm\rangle$), and $n_l = \sum_m c_{lm}^\dagger c_{lm}$ and $A_l^\dagger = (A_l)^\dagger = \sum_m c_{lm}^\dagger c_{l\bar{m}}^\dagger$ are the corresponding number- and pair-creation operators. The pairing strengths $g_{ll'}$ are here approximated by a single constant g, with ϵ_l the single-particle level corresponding to the m-fold degenerate states $|\,lm\rangle$.

As is well-known, the pairing model in Eq. (4) is central in the theory of superconductivity. Richardson's exact solution of the model [1], exploiting its integrability, has been important for applications in mesoscopic and nuclear physics where the small number of fermions prohibits the use of conventional BCS theory [4]. Moreover, its (pseudo)spin representation in the guise of the Richardson-Gaudin model, Eq. (1), provides a striking link between quantum magnetism and pairing phenomena, two central concepts in the physics of quantum matter.

The eigenstates of the Richardson-Gaudin Hamiltonian, Eq. (1), can be constructed algebraically using the quantum inverse scattering method (QISM) [5, 6]. The main objects of this method are the classical r-matrices

$$r(\lambda,\,\mu) = \frac{4}{\lambda-\mu}\sum_\alpha S^\alpha \otimes S^\alpha\Big|_{s=\frac{1}{2}} \simeq \frac{1}{\lambda-\mu}\begin{pmatrix} 1 & 0 & 0 & 0 \\ 0 & -1 & 2 & 0 \\ 0 & 2 & -1 & 0 \\ 0 & 0 & 0 & 1 \end{pmatrix}. \tag{5}$$

and the L-matrix of the loop algebra $\mathcal{L}(sl(2))$ generators $h(\lambda)$, $X^+(\lambda)$, $X^-(\lambda)$

$$L(\lambda) = \begin{pmatrix} h(\lambda) & 2X^-(\lambda) \\ 2X^+(\lambda) & -h(\lambda) \end{pmatrix}. \tag{6}$$

The commutation relations (CR) of loop algebra generators are given in compact matrix form

$$[L_1(\lambda), L_2(\mu)] = -[r_{1\,2}(\lambda, \mu), L_1(\lambda) + L_2(\mu)], \tag{7}$$

where $L_1(\lambda) = L(\lambda) \otimes \mathbb{I}$, $L_2(\mu) = \mathbb{I} \otimes L(\mu)$ and $r(\lambda,\, \mu)$ is the 4×4 c-number matrix in Eq. (5). A consequence of this form is the commutativity of transfer matrices,

$$t(\lambda) = \frac{1}{2}\mathrm{tr}_0(L^2(\lambda)) \in \mathcal{L}(sl(2)), \qquad [t(\lambda), t(\mu)] = 0. \tag{8}$$

The corresponding mutually commuting operators extracted from the decomposition of $t(\lambda)$ define a Gaudin model [2, 7]. However, to get the Richardson Hamiltonian a mild change of the L-operator is necessary,

$$L(\lambda) \to L(\lambda; c) := c\, h_0 + L(\lambda),$$

where $h_0 = \sigma_0^z$ in auxiliary space $\mathbb{C}_0^2$ of spin $1/2$. This transformation does not change the CR of matrix elements of this matrix $L(\lambda; c)$ due to the symmetry of the r-matrix (5):

$$[Y \otimes \mathbb{I} + \mathbb{I} \otimes Y, r(\lambda,\, \mu)] = 0, \quad Y \in sl(2). \tag{9}$$

The resulting transfer matrix obtains some extra terms

$$t(\lambda; c) = \frac{1}{2}\mathrm{tr}_0\left(L(\lambda; c)\right)^2 = c^2\mathbf{1} + c\, h(\lambda) + h^2(\lambda) + 2\left(X^+(\lambda)X^-(\lambda) + X^-(\lambda)X^+(\lambda)\right). \tag{10}$$

Let us consider a spin-1/2 representation on the auxiliary space $V_0 \simeq \mathbb{C}^2$ and spin ℓ_k representations on quantum spaces $V_k \simeq \mathbb{C}^{\ell_k+1}$ with extra parameters ϵ_k corresponding to site $k = 1, 2, \ldots, N$. The whole space of quantum states is $\mathcal{H} = \otimes_1^N V_k$ and the highest weight vector (highest spin, "ferromagnetic state") $|\,\Omega_+\rangle$ satisfies

$$X^+(\lambda)\,|\,\Omega_+\rangle = 0, \qquad h(\lambda)\,|\,\Omega_+\rangle = \rho(\lambda)\,|\,\Omega_+\rangle, \tag{11}$$

where

$$\rho(\lambda) = \sum_{k=1}^{N} \frac{l_k}{\lambda - \epsilon_k}.$$

It is useful to introduce the notation Y_{gl} for global operators of the $sl(2)$-representation:

$$Y_{gl} := \sum_{k=1}^{N} Y_k.$$

To find the eigenvectors and spectrum of $t(\lambda)$ on $\mathcal{H}$ one requires that vectors of the form

$$|\mu_1, \ldots, \mu_M\rangle = \prod_{j=1}^{M} X^-(\mu_j)\,|\,\Omega_+\rangle \tag{12}$$

are eigenvectors of $t(\lambda)$,

$$t(\lambda)|\{\mu_j\}_{j=1}^M\rangle = \Lambda(\lambda;\{\mu_j\}_{j=1}^M)|\{\mu_j\}_{j=1}^M\rangle, \tag{13}$$

provided that the parameters μ_j satisfy the Bethe equations:

$$2c + \sum_{k=1}^{N} \frac{\ell_k}{\mu_i - \epsilon_k} - \sum_{j\neq i}^{M} \frac{2}{\mu_i - \mu_j} = 0, \quad i = 1, \ldots, M. \tag{14}$$

The realization of the loop algebra generators on the space $\mathcal{H}$ takes the form

$$h(\lambda) = \sum_{k=1}^{N} \frac{h_k}{\lambda - \epsilon_k}, \quad X^-(\lambda) = \sum_{k=1}^{N} \frac{X_k^-}{\lambda - \epsilon_k}, \quad X^+(\lambda) = \sum_{k=1}^{N} \frac{X_k^+}{\lambda - \epsilon_k}. \tag{15}$$

The coupling constant g of (1) is connected with the parameter $c = 1/g$, while the Hamiltonian (1) is obtained as the operator coefficient of the term $1/\lambda^2$ in the expansion of $t(\lambda; c)$ at $\lambda \to \infty$.

Quantum group theory gives the possibility to deform a Hamiltonian preserving integrability [8, 9]. Specifically, we can use the so-called Jordanian r-matrix to quantum deform the Hamiltonian of Richardson-Gaudin model (1). We add to the $sl(2)$ symmetric r-matrix (5) the Jordanian part

$$r^J(\lambda,\, \mu) = \frac{C_2^{\otimes}}{\lambda - \mu} + \xi \left(h \otimes X^+ - X^+ \otimes h\right), \tag{16}$$

with Casimir element $C_2^{\otimes}$ in the tensor product of two copies of $sl(2)$,

$$C_2^{\otimes} = h \otimes h + 2\left(X^+ \otimes X^- + X^- \otimes X^+\right).$$

After the Jordanian twist the r-matrix (14) is commuting with the generator X_0^+ only,

$$[X_0^+ \otimes \mathbb{I} + \mathbb{I} \otimes X_0^+, r^{(J)}(\lambda, \mu)] = 0. \tag{17}$$

Hence, one can add the term $cX_0^+ + L(\lambda, \xi)$ to the L-operator. This yields the twisted transfer-matrix $t^{(J)}(\lambda) = \frac{1}{2}\mathrm{tr}_0(cX_0^+ + L(\lambda, \xi))^2$,

$$t^{(J)}(\lambda) = cX^+(\lambda) + h(\lambda)^2 - 2h'(\lambda) + 2(2X^-(\lambda) + \xi)X^+(\lambda). \tag{18}$$

The corresponding commutation relations between the generators of the twisted loop algebra

are explicitly given by

$$\begin{aligned}
[h(\lambda), h(\mu)] &= 2\xi \left(X^+(\lambda) - X^+(\mu)\right) \\
\left[X^-(\lambda), X^-(\mu)\right] &= -\xi \left(X^-(\lambda) - X^-(\mu)\right), \\
\left[X^+(\lambda), X^-(\mu)\right] &= -\frac{h(\lambda) - h(\mu)}{\lambda - \mu} + \xi X^+(\lambda), \\
\left[X^+(\lambda), X^+(\mu)\right] &= 0, \\
\left[h(\lambda), X^-(\mu)\right] &= 2\frac{X^-(\lambda) - X^-(\mu)}{\lambda - \mu} + \xi h(\mu), \\
\left[h(\lambda), X^+(\mu)\right] &= -2\frac{X^+(\lambda) - X^+(\mu)}{\lambda - \mu}.
\end{aligned} \tag{19}$$

The realization of the Jordanian twisted loop algebra $\mathcal{L}_J(sl(2))$ with CR (19) is given similar to (15) with extra terms proportional to the deformation parameter ξ,

$$h(\lambda) = \sum_{k=1}^{N} \left(\frac{h_k}{\lambda - \epsilon_k} + \xi X_k^+ \right), \; X^-(\lambda) = \sum_{k=1}^{N} \left(\frac{X_k^-}{\lambda - \epsilon_k} - \frac{\xi}{2} h_k \right), \; X^+(\lambda) = \sum_{k=1}^{N} \frac{X_k^+}{\lambda - \epsilon_k}. \tag{20}$$

To construct eigenstates for the twisted model one has to use operators of the form [9, 10]

$$B_M(\mu_1, ..., \mu_M) = X^-(\mu_1)(X^-(\mu_2) + \xi)...(X^-(\mu_M) + \xi(M-1)), \tag{21}$$

and acting by these operators on the ferromagnetic state $|\Omega_+\rangle$.

The deformed Richardson-Gaudin model Hamiltonian can now be extracted from the transfer-matrix $t^{(J)}(\lambda)$ as the operator coefficient in its expansion $\lambda \to \infty$.

According to Eqs. (8) and (18) one can also extract quantum integrals of motion J_k using the realization (20), reading off the expressions for J_k from the expansion

$$t^{(J)}(\lambda) = J_0 + \frac{1}{\lambda} J_1 + \frac{1}{\lambda^2} J_2 + \dots. \tag{22}$$

The corresponding quantum deformed Hamiltonian reads

$$H \simeq J_2 = c \sum_{j=1}^{N} \epsilon_j X_j^+ + 2\xi \left\{ \left(\sum_{j=1}^{N} \epsilon_j h_j \right) X_{gl}^+ - h_{gl} \sum_{j=1}^{N} \epsilon_j X_j^+ \right\} + \left({h_{gl}}^2 + 2h_{gl} + 4X_{gl}^- X_{gl}^+ \right). \tag{23}$$

It is instructive to write down a simplified case without the Jordanian twist: $\xi = 0$. One thus obtains

$$J_0 = 0, \quad J_1 = X_{gl}^+, \quad J_2 \simeq \sum_{k=1}^{N} \epsilon_k X_k^+ + \frac{g}{2} \left({h_{gl}}^2 + 2h_{gl} + 4X_{gl}^- X_{gl}^+ \right). \tag{24}$$

The case $\xi = 0$ can also be obtained by taking off from the inhomogeneous XXX spin chain. The model can be described by a 2×2 monodromy matrix [5]

$$T(\lambda) = \begin{pmatrix} A(\lambda) & B(\lambda) \\ C(\lambda) & D(\lambda) \end{pmatrix},$$

satisfying the quadratic relations

$$R(\lambda, \mu) T(\lambda) \otimes T(\mu) = (I \otimes T(\mu)) \, (T(\lambda) \otimes I) \, R(\lambda, \mu). \tag{25}$$

If we multiply $T(\lambda)$ by a constant 2×2 matrix $\mathcal{M}$ the resulting matrix $\widetilde{T}(\lambda) = \mathcal{M} \cdot T(\lambda)$ will satisfy the same relation (25). Choosing a triangular matrix

$$\mathcal{M}(\epsilon) = \begin{pmatrix} 1 & \varepsilon \\ 0 & 1 \end{pmatrix},$$

the entries of monodromy matrices become simply related:

$$\widetilde{A} = A + \varepsilon C, \quad \widetilde{B} = B + \varepsilon D, \quad \widetilde{C} = C, \quad \widetilde{D} = D.$$

This choice of $\mathcal{M}(\epsilon)$ (of the same type as considered in Ref. [11]) permits us to use the same reference state $|\Omega_+\rangle \in \mathcal{H}$ (11) and $\widetilde{B}$ as a creation operator of the algebraic Bethe ansatz [5].

Bethe states are given by the same action of product operators $\widetilde{B}(\mu_j) = B(\mu_j) + \varepsilon D(\mu_j)$ although operators $B(\mu_j)$ do not commute with $D(\mu_j)$:

$$D(\lambda) B(\mu) = \alpha(\lambda, \mu) B(\mu) D(\lambda) + \beta(\lambda, \mu) B(\lambda) D(\mu), \tag{26}$$

where

$$\alpha(\lambda, \mu) = \frac{\lambda - \mu + \eta}{\lambda - \mu}, \qquad \beta(\lambda, \mu) = -\frac{\eta}{\lambda - \mu}.$$

For a 3-magnon state one gets due to B-D ordering in Eq. (26):

$$\prod_{j=1}^{3} \widetilde{B}(\mu_j) = \prod_{j=1}^{3} B(\mu_j) + \varepsilon \sum_{s=1}^{3} \alpha(\mu_k, \mu_s) \alpha(\mu_s, \mu_l) B(\mu_k) B(\mu_l) D(\mu_s) + \\ \varepsilon^2 \sum_{s=1}^{3} \alpha(\mu_k, \mu_s) \alpha(\mu_l, \mu_s) B(\mu_s) D(\mu_k)) D(\mu_l) + \varepsilon^3 \prod_{j=1}^{3} D(\mu_j).$$

Similar formulas are valid for M-magnon states. Hence, acting on the ferromagnet state $|\Omega_+\rangle$ we obtain filtration of states with eigenvalues of $S^z : \frac{N}{2}, \frac{N}{2} - 1, \frac{N}{2} - 2, \frac{N}{2} - 3$.

More complicated deformations of the Richardson-Gaudin model can be obtained using r-matrices related to the higher rank Lie algebras [12]. The structure of the eigenstates of the transfer matrix and their entanglement properties [13] are under investigation.

Acknowledgments We would like to thank E. Damaskinsky for useful discussions. This work was supported by RFBR grants 11-01-00570-a, 12-01-00207-a, and 13-01-12405-ofi-M2 (P.K.), and by STINT grant IG2011-2028 (A.S and H.J.).

References

[1] R. W. Richardson, *Exact eigenstates of the pairing-force hamiltonian*, J. Math. Phys. **6**, 1034 - 1051 (1963).

[2] M. Gaudin, *La fonction d'onde de Bethe*, chapter 13 (Masson, Paris, 1983).

[3] M. C. Cambiaggio, A. M. F. Rivas, and M. Saraceno, *Integrability of the pairing hamiltonian*, Nucl. Phys. A **624**, 157 - 167 (1997).

[4] For a review, see J. Dukelsky, S. Pittel, and G. Sierra, *Exactly solvable Richardson-Gaudin models for many-body quantum systems*, Rev. Mod. Phys. **76**, 643 - 662 (2004).

[5] L. D. Faddeev, *How algebraic Bethe ansatz works for integrable models*, in A. Connes, K. Gawedzki, and J. Zinn-Justin, editors, *Quantum symmetries/Symetries Quantiques, Proceedings of the Les Houches summer school, session LXIV*, pages 149–219 (North-Holland, Amsterdam, 1998).

[6] P. P. Kulish and E. K. Sklyanin, *Quantum spectral transform method. Recent developments,* Volume **151** of *Lecture Notes in Physics*, pages 61–119 (Springer, New York, 1982).

[7] E. K. Sklyanin, *Generating function of correlators in the* $\mathfrak{sl}_2$ *Gaudin model,* Lett. Math. Phys. **47**, 275–292 (1999).

[8] P. P. Kulish and A. A. Stolin, *Deformed Yangians and integrable models*, Czech. J. Phys. **12**, 207–1212 (1997).

[9] P. P. Kulish, *Twisted* $sl(2)$ *Gaudin model, preprint PDMI 08/2002.*

[10] N. C. Antonio and N. Manojlovic, sl_2 *Gaudin model with Jordanian twist*, J. Math. Phys. **46**, 102701 (2005).

[11] E. Mukhin, V. Tarasov, and A. Varchenko, *The* $\mathfrak{gl}_2$ *Bethe algebra associated with nilpotent element*, arXiv:0810.2257 (2008).

[12] A. A. Stolin, *On rational solutions of Yang-Baxter equation for* $sl(n)$, Math. Scand. **69** 57 - 80 (1991).

[13] C. Dunning, J. Links, and H.-Q. Zhou, *Ground-state entanglement of the BCS model*, Phys. Rev. Lett. **94**, 227002 (2005).